No.



8200063

THIE UNINHED SHATTES OF AMIERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

Holden's Foundation Seeds, Inc.

Williams, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF eig h-teen years from the Date of this grant, subject to the payment of the required fees and periodic replenishment of viable basic by of the variety in a public repository as provided by LAW, the right to extend the others from selling the variety, ok offering it for sale, or reproducing it, or using it in producing a hybrid or different therefrom, to the extent provided by the Plant variety Protection au' 142, as amended, 7 u.s.c. 2321 et seq.)

CORN

'LH74'

Zu Ecstimony Watercot, I have hereunto sel my hand and caused the seal of the Plant Variety Protection Office to be affixed at the City of Washington

this 26th day of May in the year of our Lord one thousand nine hundred and eighty-three.

Attest

Lesgeth H. Evers Commissioner

Plant Variety Protection Office

Agricultural Marketing Service

Seritary of Agriculture

	UNITED STATES DEPARTMEN AGRICULTURA MARK	ETING SERVICE		FORM APPROVED OMB NO. 40-R3822
	LIVESTOCK, POULTRY, GRA P CICATION FOR PLANT VARIE RUCTIONS: See Reverse.	ETY PROTECTION CERTIFICATE		ompleted application form J.S.C. 553).
	TEMPORARY DESIGNATION OF	Ib. VARIETY NAME	FOR OFFICIA	AL USE ONLY
	Ex449	LH74	PV NUMBER 82	00063
2.	KIND NAME Yellow dent corn	3. GENUS AND SPECIES NAME	FILING DATE 1/20/82	TIME 12:30 A.M.
		Zea Mays	FEE RECEIVED	DATE
4 .	FAMILY NAME (BOTANICAL)	5. DATE OF DETERMINATION	\$ 500.00	1/20/82
	Gramineae	April, 1980	\$250.00	3/21/83
6.	NAME OF APPLICANT(S)	7. ADDRESS (Street and No. or R.F.D. No., Code)	City, State, and ZIP	B. TELEPHONE AREA CODE AND NUMBER
	Holden's Foundation Seeds, Inc.	Box 839, R.R. 2 Williamsburg, Iowa 523	361n	319-668-1100
9,	IF THE NAMEO APPLICANT IS NOT A	PERSON, FORM @ 10. IF INCORPORAT	ED, GIVE STATE AND	11. DATE OF INCOR-
	ORGANIZATION: <i>(Corporation, partnership,</i> Corporation	association, etc.) DATE OF INCO	RPORATION	PORATION 1968
12.	NAME AND MAILING ADDRESS OF APP ALL PAPERS:	LICANT REPRESENTATIVE(S), IF ANY, TO	SERVE IN THIS APPL	CATION AND RECEIVE
	Mr. Art L. Johnson, Bo	ox 839, Williamsburg, Id	owa 52361	
13.	CHECK BOX BELOW FOR EACH ATTAC	CHMENT SUBMITTED:		
		reeding History of the Variety (See Secti	on 52 of the Dlant Var	rights Dustration Act
	13A. Exhibit A, Origin and B.	reeding History of the Variety (See Secu	on 32 oj ine Fiant vai	iety Froiection Aci.)
	1 3B. Exhibit B, Novelty Staten	nent.		
	I. Y. Exhibit C. Objective Descript	ion of the Variety (Request form fr	COM Plant Variety Prote	ction Office
	 1	<u>-</u>	Om 1 11711 variety 1700	cuon Office.)
	☐ 130. Exhibit D, Additional Des	cription of the Variety.		
140.	DOES THE APPLICANT(s) SPECIFY THAT SEE D? (See Section 83(a). (If "Yes," and	T SEED OF THIS VARIETY BE SOLD BY V_{C} were 1.4 R ad 14 C below.) $C = I YES$	ARIETY NAME ONLY AS	A CLASS OF CERTIFIED
14b.	DOES THE APPLICANT(S) SPECIFY THA	T THIS VARIETY BE 14c, IF "YES," TO 14	B, HOW MANY GENER	ATIONS OF PRODUC-
	LIMITED AS TO NUMBER OF GENERA	TIONS? TION BEYOND	BREEDER SEED?	
	a YES 📙 NO	FOUNDATION	I REGISTERED	CERTIFIED
15a.	DID THE APPLICANT(S) FILE FOR PROTE name Of countries and dates.)	ECTION of this variety in other count i	RIES? YES	☐ × ☐ (If"Yes," give
16b.	HAVE RIGHTS BEEN GRANTED THIS V and dates.)	ARIETY IN OTHER COUNTRIES7 🗍 🕬	I H "Yes,	"give name of countries
16.	JOURNAL7 X YES	HE PUBLICATION OF HIS/HER (THEIR) NA		
17.		ble ample of basic seed of this variety will		application and will be
	The undersigned applicant(s) is (are) variety is distinct, uniform, and stable 42 of the Plant Variety Act.	ce with such regulations as may be applica the owner(s) of this sexually reproduced n au required in Section 41, and is entitled t	novel plant variety, and to protection under the	provisions of Section
		se representation herein cam jeopardize pi	orection and result in	penalties.
	DECEMBER 23 1981 NO	5 PER / 1/1	gentel Hell	He-
	DECEMBER 23, 1981 AG	TER OF 8/6/82	(SIGNATURE OF APPL	ICANT)
	(DATE)		(CICNIATURE OF ARC	LICANT
	(DATE)		(SIGNATURE OF APP	LICANT)

Exhibit A

Origin and Breeding History

LH74 is an inbred developed through a pedigreed breeding system. The originating cross was between A632 (University of Minnesota release) and B73 (Iowa State University release).

Item I is a schematic outline for the development of LH74, nursery row numbers, nursery locations, and year in which the progeny rows were grown are given.

Attached to Item I are copies of Holden's Foundation Seeds nursery books. Rows involved in the development of LH74 have been underscored.

Progeny from self-pollinated seed of LH74 are very uniform. Off-types occurring in the population have been attributed to outcrosses. These outcrosses grow more rapidly, are usually darker green, and have a larger tassel.

LH74 has been self-pollinated ear to row an adequate number of generations to be stable and uniform. Both field and ear inspections by the Iowa Crop Improvement Association in 1981 have been approved for certification of LH74, therefore indicating its uniformity of plant and ear type. Attached is a copy of those reports.

Item I LH74=Ex449=A632 \times B73=Selection I79-7812

Nursery Row No.		<u>Location</u>	<u>Year</u>
	A632 B73		
2829	A632 x B73	Iowa	1974
5560	A632 x B73 ₂₀ 1	Iowa	1975
755	A632 x B73 ₂₆ 2	Iowa	1976
1896	A632 x B73 _⊗ 3	Iowa	1977
5067	A632 x B73 ₈₂ 4	Hawaii	1977-78
5677	A632 x B73 ₈₂ 5	Iowa	1978
5605	A632 x B73 ₂₀ 6	Hawaii	1978-79
7812	A632 x B73 _Q 7	Iowa	1979

sucration, Miles, 10Wa

Cori : Sorghum

8200063

Foundation Inbred lines and/or Single Crosses

APPLICATION FOR CERTIFICATION

FIELD NO. 2/

Name of Applicant Holden Foun	lation =	Sood ne	County WhereApplicant Lives	Lowa
Address Box 839 P. 0.		et or R.R.	County Where Field is Located	Towa
Williams bur City			<i>5236 (</i> Zip Code	319 / 668 ≠ 1/0 0 Area Code/Tel ephone
Crop: (check one) Co	orn c l	Sorghum c 1	Crop Year <u>/ 9</u>	8 1
Send this completed ap Improvement Association, Am and make checks payable to is required for each field.	es, Iowa. the <u>Iowa C</u>	See "Certifica rop Improvement	tion Requirements" <u>Association</u> . A se	for amount of fee parate application
		PLANTING	SOURCE OF SEEDSTOO PLANTED*	eks
PEDI GREE POLLEN PARENT:	ACRES	<u>DATE</u>	Cert. Grown Lot No. By	REMARKS**
1. <u>LH 74</u>	12.48	April		(b)(1) H.F.5-1980 (z) Ve s
SEED PARENT:				
1. MellVGms Ht	. 7	Apr. 1 27	H.F. 5	For Insport on Only
2. Mal7 Ht	-33			- For the spectain only
3. <u>/ 1451</u>				- For thospection only
4. <u>LH58</u>		April 27	<u> </u>	- Touchage the only
5. <u>LH 58 VG ms</u>	<u>. 33</u>	•		= 70 chapeto or
b		<u> </u>		
PLANTING RATIO: SEED4/				DW WIDTH 3 a "
Check here if this				
Check here if thi *SEEDSTOCK PLANTED:	.s producti	on is by <u>Hand</u>	POLITINACION,	
(a) For Certified Seed	lstocks:	,		
			each.lot of seed pl s or pounds planted	
(b) Complete only for			s or pounds pranced	on cach cag.
(1) Originator S. (2) Does the inbrece Certification (3) If you are not	meet the Requirement the origi	definition of ots? Yes	an inbred as outline No proof that the seed rements.	

FIELD INSPECTION REPORT

Applicant #	OLDE	S.K				
Field Identity	POLLEN	74	Ch	ADD	RESS	Z /
Date 8/4	TOLLEN	TARLINI			M	P.M.
Your attention described field.	on is call	ed to the	e followin	g condit	Time ions in t	
	<u> 1</u>	POLLEN	PAREN	<u>IT</u>		
Plants shedding Off-type plants	pollen_shedding	pollen_	%. (*)	<u></u> %		Š
_		SEED	PARENT		- 4	
% Plants	とHフリ	1. E. A.	A. Lea	Siec /	40/1/2	102
Shedding Pollen	100	٥	ဂ	0	٥	0
Receptive to Pollen	954					
Off-type		۵	0	0	ဂ	0
Off-type & Shed- ding Pollen	D	0_	0	0	o o	0
REMARKS:						
	FIN	AL	125	1077	01)	
FIELD	LOOK	5 0	000	15	01 (<u> ッド </u>
					4/	7/
				12	tall	<u>/</u>
GROWER'S COPY				7	nspector [3-79-10M

5+12 No 21

Pollmator LH 74- Early

	Dote	Dryer	Bin	Podigree	Bushols	Inspection Results	Tasportor
1		3	30-29-28 27-26-25 24-23-22	1 4 74 Blook	650		210
	9/21		24-23-22	1 14 17 OWNE		Bassed 9/21/81	AL
ł		 	 			7	
1			1				
į į					1		
ŀ		 		· · · · · · · · · · · · · · · · · · ·			
ŀ		 	 				
ŀ		ļ					<u> </u>
}			1				
ŀ							ļ
	Dote	Dryor	Bin	Padigica	Bushols	Inspection Results	JASpecto,
. ↓							
			,				
Ī		1	-				
ļ							
.							
ľ		-					
ŀ							
ŀ		 					+
ł		ļ 					
ŀ	<u>. </u>	<u></u>		· · · · · · · · · · · · · · · · · · ·			+
ł							+
ı							
ŀ		 	<u> </u>				
ļ			<u> </u>				
ļ	,						
ļ		1		_			<u> </u>
ŀ	·						
ļ							
Ī							
Ī							
1							
Ī							-
ł							
- 1			 				 -

Supplement to Exhibit A

Corn Application No. 8200063, 'LH74'

LH74 is very stable and uniform. Either self-pollination or sib mating will maintain the identity of LH74 very well.

LH74 has been increased three generations since the line development process was completed in 1979. Each increase generation is carefully scrutinized for uniformity.

Supplement to Exhibit A

Corn Application #8200063 'LH74'

There are no variants in LH74 inbred.

-

Exhibit B

LH74 most closely resembles A632Ht; however, at flowering A632Ht's leaves are horizontal whereas LH74's leaves are upright with only the end one-third of the leaf broken over.

Photo 1 illustrates LH74's upright leaf, and Photo 2 shows A632Ht's flat leaf angle. Photo 3 compares A632Ht on the left and LH74 on the right.



Photo 1








Photo 2

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE LIVESTOCK, POULTRY, GRAIN & SEED DIVISION BELTSVILLE, MARY LAND 20706

OBJECTIVE DESCRIPTION OF VARIETY

CORN (ZEA MA YS)

NAME OF APPLICANT(S)	EOD OFFICIAL LISE AND V			
Holden's Foundation Seeds, Inc.	P VPO NUMBER COOLOGO			
	8200063			
	DESIGNATION			
Williamsburg, Iowa 52361	1. H 74			
Place the appropriate number that describes the varietal character of this variety in the Place a zero in first box (**** 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	boxes below.			
	7 - 1033			
1 -SWEET 2 = DENT 3 = FLINT 4 = FLOUR 5 = F	OP 6 ∓ ORNAMENTAL			
2. REGION WHERE BEST ADAPTED IN THE U.S.A.:				
2 1 NORTHWEST 2 NORTHCENTRAL 3 NORTHEAST c 1 6 SOUTHCENTRAL 6 SOUTHWEST 7 MOST REGIONS	4 -SOUTHEAST			
	comments" (pg. 3) state how			
heut uni	ts were calculated)			
6 1 days from emergence to 60% of plants in silk	4 N4 I T s			
DAYS FROM 50% SILK TO OPTIMUM EDIBLE QUALITY	TUNITS			
8 3 ave soon for the majority of the majority of				
Holden's Foundation Seeds, Inc. **POPOLINES** (Stream of the green for the content of the variety in d c bases below. Place a zero in first bas (**+ 3 **) for 0 **) when number is either 99 or less or 9 or less. **Place a zero in first bas (**+ 3 **) for 0 **) when number is either 99 or less or 9 or less. **Place a zero in first bas (**+ 3 **) for 0 **) when number is either 99 or less or 9 or less. **I TYPE: **2				
4. PLANT:				
1 9 0	CM FAR HEIGHT (To base of ton 881)			
1 8 9 CM, HEIGHT (TO tasset tip)	1/2 CIVIL EAR HEIGHT (10 Base of top said			
BOX 839 Williamsburg, Iowa 52361 Place the appropriate number that describes the varietal character of this variety in (i c bases below, Place a zero in first box (*4. 0 19 9 or 0.9) when number is either 99 or less or 9 or less. 1. TYPE: 2 1 **SWEET 2 **DENT 3 **FIRST 4 **SOUTHEAST 4 **SOUTHEAST 2 **NORTHEAST 4 **SOUTHEAST 7 **MOST REGIONS 2 ** MOST REGIONS 2 ** MOST REGIONS 2 ** MOST REGIONS 3 ** MATURITY (in Region of Best Adaptability): 6 1 DAYS FROM EMERGENCE TO 40% OF PLANTS IN SILK DAYS FROM 50% SILK TO OPTIMUM EDIBLE QUALITY (FA) 3 ** DAYS FROM 50% SILK TO OPTIMUM EDIBLE QUALITY (FA) 4 ** PLANT: 1 ** SUBJECT 2 ** DENT 3 ** PLANTS IN SILK DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE 1 2 ** NORTHEAST 1 ** SOUTHEAST 1 ** SUBJECT CONTROL OF TOP EAR INTERNODE **NORTHEAST 4 ** SOUTHEAST 4 ** SOUTHEAST 1 ** SOUTHEAST 5 ** MOST REGIONS 50% SILK TO OPTIMUM EDIBLE QUALITY (FB, 3) ** SIZE HOW heat strift were ediculated (FA) 3 ** A STRUM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE 1 2 ** TILL TO SIZE HOW HEAT TO SIZE				
Holden's Foundation Seeds, Inc. ADDRES (Greet and to by EFD. No. City, Mate, and APP Code) Box 839 Williamsburg, Iowa 52361 Place the appropriate number that describes the varietal character of this variety in (1 c bases below. Place 2 fot in first bas (*** [0] 9 9 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0				
Number of Tillers: Number of Ears Par Stalk	:			
2 NONE 2-1-2 $3 = 2-3$ $4 = > 3$ $c/3$ 1-SINGLE 2	-SLIGHT TWO-EAR TENDENCY			
2 1 NONE 2-1-2 3=2-3 4=>3 C/31-SINGLE 2	-SLIGHT TWO-EAR TENDENCY			
2) NONE 2-1-2 3=2-3 4=>3 Cl 31-SINGLE 2 3 STRONG TWO	SLIGHT TWO-EAR TENDENCY -EAR TENDENCY 4 = THREE-EAR TENDENCY			
2) NONE 2-1-2 3=2-3 4=>3 Cl 31-SINGLE 2 3 STRONG TWO	SLIGHT TWO-EAR TENDENCY -EAR TENDENCY 4 = THREE-EAR TENDENCY			
2 I NONE 2-1-2 3=2-3 4 > 3 C/31-SINGLE 2 3 STRONG TWO Cytoplasm Type: 1= NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER	SLIGHT TWO-EAR TENDENCY -EAR TENDENCY 4 = THREE-EAR TENDENCY			
2) NONE 2-1-2 3 = 2-3 4 = >3 C/31-SINGLE 2 3 = STRONG TWO Cytoplasm Type: 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5, LEAF (Field Corn Inbred Examples Given);	SLIGHT TWO-EAR TENDENCY -EAR TENDENCY 4 = THREE-EAR TENDENCY			
2 NONE 2-1-2 3 = 2-3 4 = >3 C/31-SINGLE 2 3 STRONG TWO Cytoplasm Type: 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 OTHER 5. LEAF (Field Corn Inbred Examples Given); Color: 1 = LIGHT GREEN (HY) 2 MEDIUM GREEN (WF9) 3 DARK GR	SIGHT TWO-EAR TENDENCY -EAR TENDENCY 4 THREE-EAR TENDENCY (Specify)			
2 NONE 2-1-2 3 = 2-3 4 = >3 C/31-SINGLE 2 3 STRONG TWO Cytoplasm Type: 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 OTHER 5. LEAF (Field Corn Inbred Examples Given); Color: 1 = LIGHT GREEN (HY) 2 MEDIUM GREEN (WF9) 3 DARK GR	SIGHT TWO-EAR TENDENCY -EAR TENDENCY 4 THREE-EAR TENDENCY (Specify)			
2 I NONE 2-1-2 3=2-3 4=>3 C/31-SINGLE 2 3 II STRONG TWO Cytoplasm Type: 1 = NORMAL 2="T" 3= "S" 4= "C" 5 II OTHER 5. LEAF (Field Corn Inbred Examples Given); Color: 1 = LIGHT GREEN (HY) 2 II MEDIUM GREEN (WF9) 3 II DARK GR	SIGHT TWO-EAR TENDENCY -EAR TENDENCY 4 THREE-EAR TENDENCY (Specify)			
2 I NONE 2-1-2 3=2-3 4=>3 C/31-SINGLE 2 3 STRONG TWO Cytoplasm Type: 1 = NORMAL 2="T" 3="S" 4="C" 5 TOTHER 5. LEAF (Field Corn Inbred Examples Given); Color: 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GR c ² I Angle from Stalk (Upper half): Sheath Pubscence:	SE-SLIGHT TWO-EAR TENDENCY -EAR TENDENCY 4 THREE-EAR TENDENCY (Specify) EEN (B14) 4 VERY DARK GREEN (K166)			
2 NONE 2-1-2 3 = 2-3 4 = >3 C/31-SINGLE 2 3 STRONG TWO Cytoplasm Type: 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 OTHER 5. LEAF (Field Corn Inbred Examples Given); Color: 1 = LIGHT GREEN (HY) 2 MEDIUM GREEN (WF9) 3 DARK GR c^2 Angle from Stalk (Upper half); Sheath Pubscence: 1 = < 30° 2 30-60° 3 => 60°	SE-SLIGHT TWO-EAR TENDENCY -EAR TENDENCY 4 THREE-EAR TENDENCY (Specify) EEN (B14) 4 VERY DARK GREEN (K166) (W22) 2 MEDIUM (WF9)			
2 NONE 2-1-2 3 = 2-3 4 = >3	SE-SLIGHT TWO-EAR TENDENCY -EAR TENDENCY 4 THREE-EAR TENDENCY (Specify) EEN (B14) 4 VERY DARK GREEN (K166) (W22) 2 MEDIUM (WF9)			
2 NONE 2-1-2 3 = 2-3 4 = >3	SE-SLIGHT TWO-EAR TENDENCY -EAR TENDENCY 4 THREE-EAR TENDENCY (Specify) EEN (B14) 4 VERY DARK GREEN (K166) (W22) 2 MEDIUM (WF9)			
2 NONE 2-1-2 3 = 2-3 4 = > 3 C 3 -SINGLE 2 3 STRONG TWO Cytoplasm Type: 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 OTHER 5. LEAF (Field Corn Inbred Examples Given): Color: 1 = LIGHT GREEN (HY) 2 MEDIUM GREEN (WF9) 3 DARK GR c ² Angle from Stalk (Upper half): Sheath Pubscence: 1 = < 30	SLIGHT TWO-EAR TENDENCY -EAR TENDENCY 4 = THREE-EAR TENDENCY (Specify) EEN (814) 4 = VERY DARK GREEN (K166) (W22) 2 = MEDIUM (WF9) (OH26) (OH51) 2 = FEW (OH56A)			
2 NONE 2-1-2 3 = 2-3 4 = >3 C/31-SINGLE 2 3 = STRONG TWO Cytoplasm Type: 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given); Color: 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GR c^2 Angle from Stalk (Upper half); Sheath Pubscence: 1 1 = < 30° 2 = 30-60° 3 => 60°	SLIGHT TWO-EAR TENDENCY -EAR TENDENCY 4 = THREE-EAR TENDENCY (Specify) EEN (814) 4 = VERY DARK GREEN (K166) (W22) 2 = MEDIUM (WF9) (OH26) (OH51) 2 = FEW (OH56A)			
2 NONE 2-1-2 3 = 2-3 4 = >3 C/31-SINGLE 2 3 = STRONG TWO Cytoplasm Type: 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5. LEAF (Field Corn Inbred Examples Given); Color: 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GR c^2 Angle from Stalk (Upper half); Sheath Pubscence: 1 1 = < 30° 2 = 30-60° 3 => 60°	SLIGHT TWO-EAR TENDENCY -EAR TENDENCY 4 = THREE-EAR TENDENCY (Specify) EEN (814) 4 = VERY DARK GREEN (K166) (W22) 2 = MEDIUM (WF9) (OH26) (OH51) 2 = FEW (OH56A)			
2 NONE 2-1-2 3=2-3 4=>3 Cl 31-SINGLE 2 3 STRONG TWO Cytoplasm Type: 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5, LEAF (Field Corn Inbred Examples Given); Color: 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GR c² Angle from Stalk (Upper half); Sheath Pubscence: 1 = LIGHT 3 = HEAV' Merginal Waves: CI 3 1 = NONE (HY) 2 = FEW (WF9) 3 = MANY (OH7L) 2 1 = ABSENT C 1 3 = MANY Width: Length:	SLIGHT TWO-EAR TENDENCY -EAR TENDENCY 4 THREE-EAR TENDENCY (Specify) EEN (814) 4 VERY DARK GREEN (K166) (W22) 2 MEDIUM (WF9) (OH26) (OH51) 2 FEW (OH56A) (PA11)			
2 NONE 2-1-2 3=2-3 4=>3 Cl 31-SINGLE 2 3 STRONG TWO Cytoplasm Type: 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5, LEAF (Field Corn Inbred Examples Given); Color: 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GR c² Angle from Stalk (Upper half); Sheath Pubscence: 1 = LIGHT 3 = HEAV' Merginal Waves: CI 3 1 = NONE (HY) 2 = FEW (WF9) 3 = MANY (OH7L) 2 1 = ABSENT C 1 3 = MANY Width: Length:	SLIGHT TWO-EAR TENDENCY -EAR TENDENCY 4 THREE-EAR TENDENCY (Specify) EEN (814) 4 VERY DARK GREEN (K166) (W22) 2 MEDIUM (WF9) (OH26) (OH51) 2 FEW (OH56A) (PA11)			
2 NONE 2-1-2 3=2-3 4=>3 Cl 31-SINGLE 2 3 STRONG TWO Cytoplasm Type: 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER 5, LEAF (Field Corn Inbred Examples Given); Color: 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GR c² Angle from Stalk (Upper half); Sheath Pubscence: 1 = LIGHT 3 = HEAV' Merginal Waves: CI 3 1 = NONE (HY) 2 = FEW (WF9) 3 = MANY (OH7L) 2 1 = ABSENT C 1 3 = MANY Width: Length:	SLIGHT TWO-EAR TENDENCY -EAR TENDENCY 4 THREE-EAR TENDENCY (Specify) EEN (814) 4 VERY DARK GREEN (K166) (W22) 2 MEDIUM (WF9) (OH26) (OH51) 2 FEW (OH56A) (PA11)			

TASSEL: 0 7 NUMBER OF LATERAL BRANCHES Branch Angle from Central Spike: Penduncie Length: 1 = < 309 CM. FROM TOP LEAF TO BASAL BRANCHES : 2 C I Pollen Shed: 1 = LIGHT (WF9) 2 ■ MEDIUM = YELLOW 2 = PINK 3 = RED

OTHER (Specify) 4 PURPLE 5 • GREEN OTHER (Specify) Pollen Restoration for Cytoplasms (o = Not Tested, 1 = Partial, 2 = Good) $\,$ OTHER (Specify Cytoplasm and degrees of restoration) $_{\text{C}}$ 7. EAR (Husked Ear Data Except When Stated Otherwise): MM. MID-POINT DIAMETER CM LENGTH GM. WEIGHT Kernel Rows: NUMBER c²I 1 INDISTINCT 2 = DISTINCT 1 = STRAIGHT 2 = SLIGHTLY CURVED c 1 Silk Color (Exposed at Silking Stage): 1 = GREEN 2 = PINK 3 = SALMON 4 = RED Husk Color: **FRESH** 2 = DARK GREEN 3 = PINK c^2 1 = LIGHT GREEN 4 = RED 5 = PURPLE 6 = BUFF _a6 DRYHusk Extmtion: (Harvest Stage) 1 **≠** SHORT (Ears **Exposed)** 2 **■** MEDIUM (Barely **Covering** Ear) 1 = SHORT (< 8 CM) 2 = MEDIUM (8-15 CM) 2 1 3 = LONG (8-10CM Beyond Ear Tip) $_{\rm c}^2$, 3 = LONG (> 15 CM) 4 - VERY LONG (> 10 CM) Position at Drv Husk Stags: cM LONG NO. OF INTERNODES $_{\rm c}1$ ₁ = UPRIGHT 2 = HORIZONTAL Taper: Drying Time (Unhusked Ear): 3 1=SLOW 2 = AVERAGE 1 = SLIGHT 3 = EXTREME 2 = AVERAGE 3 **≖** FAST KERNEL (Dried): Site (From Ear Mid-Point): MM LONG MM. WIDE MM. THICK Shape Grade (% Rounds) а 5 4 = 60-80 1 * < 20 2 = 20-40 3 = 40-60 5 = > 80

8 KERNEL	(Dried) :						8200	0063
1	Pericarp color:	1 = COLORLESS 5 = BROWN 8 = BADEGATED (6 - 'L	RED-WHITE		3 = TAN 7 = CHERRY		
	Aleurone Color:	1 = HOMOZYGOUS	2	2 ■ SEGREC	ATING (Describe) _			
1	•	2 ■ PINK 8 = PALE PURPL			BROWN GATED (Describe)		5 = BRONZE	6 × R€C
3 3	Endosperm Color:	1 ■ WHITE 2	■PALE Y	ELLOW 3	■ YELLOW	4 = PINK-	ORANGE 5 = WHI	TE CAP.
Endosper	m Type:							
a 3	1 SWEET ((su1) 2 = EXTRA CH 6 = HIGH			3 ■ NORMAL ST 7 = HIGH LYSINE		HIGH AMYLOSE ST OTHER (Specify)	ARCH
2 4	GM, WEIGHT /10	O SEEDS (Unsized Sam	pie)					
9.1508 : 2 9	MM. DIAMETER	AT MID-POINT						
ngth;	I-WEAK	2 =	STRONG	Color:	1 ■ ■ WHITE 2		RED 4 BROWN R (Specify)	
10. DISEASE	RESISTANCE (0 =	Not Tested. 1 = Susce	eptible, 2 = R	esistant);				
	STALK ROT (Dig	ata dia l	STALK	DOT 15	atu. N	0	CTALK DOT 15th	
						а	STALK ROT (Gibbere	lla)
0		LEAF BLIGHT			BLIGHT	0	SMUT	
لاـــ		ust 1 0	1	SMUT		<u>c </u>	BACTERIAL WILT	
0		L	0 MAIZE	DWARF N	MOSAIC	0	STUNT	
I a	OTHER (Specify)							
11. INSECT	RESISTANCT (O =	Not Tested, 1 ■ Susce	eptible2 = Res	sistant):				
0	CORNBORER	a 0 EAI	RWORM		0 SAP	BEETLE	О АРНІ	D
لما	ROOTWORM (No	a^{0} RO	OTWORM (W	estern)				
c^0 l	ROOTWORM (Se	outhern) $egin{array}{c} 0 & OTF \end{array}$	HER (Specify)				
12. VARIETII	ES MOST CLOSE	L Y RESEMBLING THAT.	SHRMITTEE	<u>Σ ΕΘΦ Τ<i>ΙΙ</i></u>	ℯℰℹ⅄℩ⅈ℩⅄ ℴ⅂℟ℸ℞ℴℷ⅀℣	VEN:		
CHARAC	TER		/ARIFTY	<u> </u>	CHARA <u>C</u> TER		VARIETY	
aturit <u>y</u>			32 <u>Ht</u>	- 11	Kernel Type		A632Ht	
Plant Ty			3Ht 32Ht		Quality (Edibl	<u>e)</u>	A632Ht	
Ear Type		/ 1101	72110	<u> </u>	Usage			
REFEREN	U.S. Department Corn: Culture, Pro Emerson. R.A., G.V. Tha Mutants of M Stringfield, G.H.	Agriculture. Yearbook Docessing, Products. 1976 N. Beadle, and A.C. Fras Bize, 1968. Crop Science Walze Inbred Lines of Company for the City	O Avi Publish ser. A Summe a Saclety of Ohlo. Ohlo A.	America. M E.S. Bui, 83	ge Studies in Malze.C adison, Wisconsin. 1. 1959.	Cornell A.E.S.,	Mem. 190. 1935	
		= A System for the Cla	SSITICATION OF	COLL HIDLED	Lines i riių. Inesis,	, Jino State U	iiversity,	
COMME	N15:	Hi	Temp <	4 86"	F + Low Te	mp > 50	" F	
	Heat	Units= —				_ =_	 -50	

Exhibit D

LH74 very greatly resembles A632Ht. LH74 will pollinate at the same time as A632Ht; however, LH74 most generally has silks present before pollen is shedding. Therefore, the silk date is 3-4 days earlier than the silk date of A632Ht.

LH74's plant height is only a few centimeters taller than A632Ht's. Ear height is the same as A632Ht.

Open pollinated seed set of LH74 is much better than that of $A\,6\,3\,2H\,\text{t}\,\text{.}$

LH74 is darker green than A632Ht and comes closer to the color of B73Ht.

The tassel of LH74 is smaller than that of A632Ht. The size generally resembles B73Ht, but the branches relax at anthesis like A632Ht whereas the branches of B73Ht stay erect.

The leaf angle of LH74 is most like that of B73Ht.